

# PROBING NEW PHYSICS IN $B_s$ AND $D$ MIXINGS, AND $A_{CP}(B^+ \rightarrow J/\psi K^+)$

GEORGE W.S. HOU\*

*Department of Physics, National Taiwan University, Taipei, Taiwan 10617, R.O.C.*

*\*E-mail: wshou@phys.ntu.edu.tw*

A 4th generation could be consistent with the recently measured  $\Delta m_{B_s}$  as well as  $\mathcal{B}(b \rightarrow s\ell^+\ell^-)$ , which are SM-like, but generate large  $\sin 2\Phi_{B_s} \simeq -0.5$  to  $-0.7$ . The sign is determined by the hint for New Physics in CPV measurements in charmless  $B$  decays. The  $4 \times 4$  unitarity allows one to connect to all processes involving flavor. Fixing  $V_{t'b}$ ,  $V_{t's}$  and  $V_{t'd}$  by  $Z \rightarrow b\bar{b}$ ,  $b \rightarrow s$  and  $s \rightarrow d$  processes, we predict  $D$  mixing to be close to the current bound. As a further corollary, we suggest that  $A_{CP}(B^+ \rightarrow J/\psi K^+)$  could be at 1% level or higher, where we give plausibility of an associated strong phase. Our predictions can be tested in the near future.

## 1. Introduction: SM Reigns?

The New York Times reported on July 4th the measurement of  $B_s$  mixing at the Tevatron, stating that “it was right on the money as predicted by the Standard Model”, and quoting a CDF spokeswoman, “Our real hope was for something bizarre”.

The measured<sup>1</sup>  $\Delta m_{B_s} = 17.77 \pm 0.10 \pm 0.07 \text{ ps}^{-1}$  is indeed consistent with SM, but there is still hope for something bizarre: Can CP violation in  $B_s$  mixing be large? Given that  $\sin 2\Phi_{B_s}^{\text{SM}} = -\sin 2\beta_s \sim -0.04$  is very small, *any definite measurement at the Tevatron would imply New Physics* (NP). There is reason for hope. The  $\Delta m_{B_s}$  value is somewhat lower than the CKM/UT fit projections made without using  $\Delta m_{B_s}$  in the fit.

In the 4 generation model we predict  $\sin 2\Phi_{B_s}$  is *large and negative*, with two corollaries. One is finite  $D$  mixing close to current bounds, the other is observable direct CPV (DCPV) in  $B^+ \rightarrow J/\psi K^+$  decay. Mixing dependent CPV (TCPV) measured in  $B^0 \rightarrow J/\psi K^0$ , namely<sup>2</sup>  $\mathcal{S}_{J/\psi K} = 0.685 \pm 0.032$ , is also low against CKM/UT fit predictions, which could be due to NP phase.

Admittedly, SM4 has troubles with precision EW tests.<sup>3</sup> But with the LHC approaching, we should keep an open mind. For  $N_\nu$  counting, as discussed by Soddu,<sup>4</sup> mas-

sive neutrinos call for NP. The reason we focus on the 4th generation is its ease in affecting heavy meson mixings and other electroweak penguins (EWP),<sup>5</sup> and it naturally brings in a new CPV phase.<sup>6</sup>

## 2. Large CPV in $B_s$ Mixing

The 4 generation unitarity for  $b \rightarrow s$  transitions is  $\lambda_u + \lambda_c + \lambda_t + \lambda_{t'} = 0$ , where  $\lambda_i \equiv V_{is}^* V_{ib}$ . Since  $|\lambda_u| < 10^{-3}$  by direct measurement, one effectively has

$$\lambda_t \cong -\lambda_c - \lambda_{t'}, \quad (1)$$

where one has a NP CPV phase<sup>6</sup> through  $\lambda_{t'} \equiv V_{t's}^* V_{t'b} \equiv r_{sb} e^{i\phi_{sb}}$ , and Eq. (1) becomes a triangle with potentially large area, i.e. large CPV effect.

The formula for  $B_s$  mixing is<sup>7</sup>

$$M_{12} \propto f_{B_s}^2 B_{B_s} \left\{ \lambda_c^2 S_0(t, t) - 2\lambda_c \lambda_{t'} \Delta S_0^{(1)} + \lambda_{t'}^2 \Delta S_0^{(2)} \right\}, \quad (2)$$

where  $S_0(t, t)$  gives SM3 top effect, and the  $t'$  effects are GIM subtracted and vanish with  $\lambda_{t'}$ , analogous to  $\Delta C_i \equiv C_i^{t'} - C_i^t$  terms<sup>6</sup> that modify the Wilson coefficients  $C_i$  for  $b \rightarrow s$  decays. One also has analogous strong dependence on  $m_{t'}$  is, i.e. nondecoupling of  $t$  and  $t'$  from box and  $Z$  penguins.<sup>5</sup>




Fig. 1. (a)  $\Delta m_{B_s}$ , (b)  $\sin 2\Phi_{B_s}$  vs  $\phi_{sb}$ , for  $m_{t'} = 300$  GeV and  $r_{sb} = 0.02, 0.025$  and  $0.03$ . Larger  $r_{sb}$  gives stronger variation.

Taking  $m_t = 170$  GeV and the central value of  $f_{B_s}\sqrt{B_{B_s}} = 295 \pm 32$  MeV from lattice, we find  $\Delta m_{B_s}^{\text{SM}} \sim 24$  ps $^{-1}$ , which is on the high side compared with Eq. (1). Of course  $f_{B_s}\sqrt{B_{B_s}}$  could be lower, but *it could also be higher*. One may therefore need SM4 to bring  $\Delta m_{B_s}$  down a bit.

Keeping  $f_{B_s}\sqrt{B_{B_s}} = 295$  MeV, in Fig. 1(a) we plot  $\Delta m_{B_s}$  vs  $\phi_{sb}$  for  $m_{t'} = 300$  GeV and  $r_{sb} = 0.02, 0.025$  and  $0.03$ , where dashed line is the SM3 value, and solid band is the  $2\sigma$  range of Eq. (1). We see that<sup>7</sup>  $\Delta m_{B_s}$  comes down to the CDF range in 1st and 4th quadrant. For  $r_{sb} = 0.02, 0.025, 0.03$ , we find  $\phi_{sb} \simeq 52^\circ\text{--}55^\circ, 62^\circ\text{--}64^\circ, 67^\circ\text{--}69^\circ$ . This implies large CPV, i.e. large  $\sin 2\Phi_{B_s}$ , which is plotted in Fig. 1(b).

It is important to note that the parameter range above not only gives SM-like  $\Delta m_{B_s}$ , it also gives SM-like  $\mathcal{B}(b \rightarrow s\ell^+\ell^-)$ ,<sup>8</sup> as the latter is also dominated by EWP and box diagrams.<sup>5</sup> In fact, combining  $\Delta m_{B_s}$  with the  $b \rightarrow s\ell^+\ell^-$  rate,  $|\phi_{sb}| \gtrsim 55^\circ$  is implied,<sup>7</sup> which practically rules out the allowed range from  $\Delta m_{B_s}$  for  $r_{sb} \sim 0.02$ . This leads to  $|\sin 2\Phi_{B_s}| \sim 0.5$  to  $0.7$ . Thus, things may still turn “bizarre”. Given that CDF has made precision measurement<sup>1</sup> of  $\Delta m_{B_s}$ , *can one pull off another coup in measuring*

$\sin 2\Phi_{B_s}$ , *before LHC start?* Any definite measurement would be a discovery of NP!

Currently we have two hints for NP in CPV  $b \rightarrow s$  transitions. Interestingly, they favor  $\sin 2\Phi_{B_s} < 0$ . One hint is TCPV in  $b \rightarrow s\bar{q}q$ : the  $\Delta\mathcal{S} \equiv \mathcal{S}_{s\bar{q}q} - \mathcal{S}_{\bar{c}cs} < 0$  problem.<sup>2</sup> The other hint is difference in DCPV between  $B \rightarrow K^+\pi^-$  vs  $K^+\pi^0$ : the  $-\Delta\mathcal{A}_{K\pi} \equiv \mathcal{A}_{K\pi} - \mathcal{A}_{K\pi^0} < 0$  problem.<sup>9</sup>

All measurements of TCPV in  $b \rightarrow s\bar{q}q$  modes at present give values lower<sup>2</sup> than charmonium modes, giving a combined significance of  $2.5\sigma$ . What aggravates this is the SM expectation of  $\Delta\mathcal{S} > 0$ . In QCDF, it was shown<sup>10</sup> that  $\mathcal{S}_{K\pi}$  and  $\mathcal{S}_{\phi K}$  are more robust than rates, which have large hadronic uncertainties. However,  $\mathcal{S}_{\eta'K}$  gets diluted away by effect of the large rate. In a model independent way,<sup>11</sup> it has recently been shown that, if this discrepancy persists as data improves, it would definitely imply NP.

The difference  $\Delta\mathcal{A}_{K\pi} \simeq 0.15$  is now established.<sup>9</sup> It is a puzzle because naively one expects it to be smaller. There are two possibilities. One is an enhancement of the color-suppressed tree ( $C$ ). The other is from  $P_{\text{EW}}$  (the EWP), which would demand NP CPV effect. The latter case was demonstrated with the 4th generation,<sup>8</sup> where the  $\phi_{sb}$  phase of Eq. (2) affects  $P_{\text{EW}}$ . The  $C$  and  $P_{\text{EW}}$  efforts were recently joined<sup>12</sup> and carried to NLO in PQCD factorization. Both trends for  $\Delta\mathcal{A}_{K\pi}$  and  $\Delta\mathcal{S}$  can be accounted. Interestingly, predictions for  $\mathcal{A}_{K^0\pi^0}$  and  $R_c, R_n$  ratios are in good agreement with the new experimental results,<sup>9</sup> while further prediction for  $\mathcal{S}_{\rho K}$  can be tested in the future.

As these are CPV measurables, the upshot from the  $\Delta\mathcal{S}$  and  $\Delta\mathcal{A}_{K\pi}$  discussion is that they *select*  $\sin 2\Phi_{B_s} < 0$  in SM4.

### 3. D Mixing Prediction

Four generation unitarity links all flavor changing and CPV processes together. With  $V_{t's}^*V_{t'b}$  large, one has to check for

consistency<sup>13</sup> with other processes. A typical  $4 \times 4$  CKM matrix is given in Ref. 13. One first saturates the  $Z \rightarrow b\bar{b}$  bound with  $V_{t'b} \simeq -0.22$ , which then fixes  $V_{t's}$  by  $b \rightarrow s$  effects. Applying the stringent kaon physics constraints fixes  $V_{t'd}$ .

A very important test is  $b \rightarrow d$  transitions. Remarkably, when the above procedure is done, it was found that  $B_d$  mixing and associated CPV (“ $\sin 2\phi_1$ ”), as well as other  $b \rightarrow d$  effects, all do not get much affected. The reason is because one cannot easily tell apart (at present level of errors) the  $b \rightarrow d$  unitarity quadrangle in SM4, from the triangle in SM3.

One striking feature of the “fitted”  $4 \times 4$  matrix is that  $V_{t'd} \simeq -0.0044 e^{-i10^\circ}$ ,  $V_{t's} \simeq -0.114 e^{-i70^\circ}$ , and  $V_{ub'} \simeq 0.068 e^{i61^\circ}$ ,  $V_{cb'} \simeq 0.116 e^{i66^\circ}$  are not smaller than 3rd generation elements. Though somewhat uncomfortable, this is data driven, and draws our interest to  $D$  mixing, since

$$V_{ub'}V_{cb'}^* \equiv r_{uc} e^{-i\phi_{uc}} = +0.0033 e^{-i5^\circ}, \quad (3)$$

would affect  $c \rightarrow u$  transitions via  $b'$  loops. Since  $|V_{ub}V_{cb}| \lesssim 10^{-4}$  by direct measurement, the unitarity condition is effectively

$$V_{ud}V_{cd}^* + V_{us}V_{cs}^* + V_{ub'}V_{cb'}^* \simeq 0, \quad (4)$$

with  $V_{ud}V_{cd}^* \simeq -0.218$  and  $V_{us}V_{cs}^* \simeq 0.215$  real to better than 3 decimal places, much like in SM. These govern  $c \rightarrow u\bar{d}$  and  $u\bar{s}$  processes, where especially the latter could generate width difference  $y_D = \Delta\Gamma_D/2\Gamma_D$  through long-distance effects.

Though small,  $V_{ub'}V_{cb'}^*$  of Eq. (3) can affect  $D^0\text{--}\bar{D}^0$  mixing, because  $m_{b'} \sim m_{t'}$  is expected, hence very heavy. The short distance effect corresponds to the  $\Delta S_0^{(2)}$  term in Eq. (2), with  $f_{B_s}^2 B_{B_s} \lambda_{t'} \rightarrow f_D^2 B_D |V_{ub'}V_{cb'}^*|$ , and  $m_{t'} \rightarrow m_{b'}$ . This generates  $x_D^{\text{SD}}$  which would be vanishingly small in SM3 because of  $|V_{ub}V_{cb}|^2$  suppression.

We used<sup>13</sup>  $V_{t'd}^*V_{t'b} \equiv r_{db} e^{i\phi_{db}}$  to fit kaon data, and found  $\phi_{db} \sim 10^\circ$  and  $r_{db} \sim 10^{-3}$ . For illustration, we take  $f_D\sqrt{B_D} = 200$  MeV




Fig. 2.  $\Delta m_D$  vs  $\phi_{db}$  ( $\sim 10^\circ$  expected) for  $m_{b'} = 230, 270$  and  $310$  (highest) GeV and  $r_{db} = 10^{-3}$ . The horizontal line is the experimental bound.

and plot  $\Delta m_D$  vs  $\phi_{db}$  in Fig. 2, for  $m_{b'} = 230, 270$  and  $310$  GeV. Our scenario predicts  $x_D^{\text{SD}} \equiv \Delta m_D^{\text{SD}}/\Gamma_D \sim 1\% - 3\%$ , which lies just below the current bound<sup>14</sup> (horizontal line), and could be accessible soon. We find CPV in  $D^0$  mixing to be no more than  $-0.2$  level, which is consistent with null search for CPV.

There is in fact a hint for width difference. Averaging over  $D^0$  decays to CP eigenstates  $K^+K^-$  and  $\pi^+\pi^-$  gives<sup>14</sup>  $y_{\text{CP}} = 0.90 \pm 0.42\%$ . Another effort is to measure  $x' = x_D \cos \delta + y_D \sin \delta$  and  $y' = y_D \cos \delta - x_D \sin \delta$  in wrong-sign  $D^0 \rightarrow K^+\pi^-$  decays, which could arise through mixing, or from doubly Cabibbo suppressed decays. The current best limit comes from Belle,<sup>14</sup>  $|x'| < 2.7\%$  and  $-1\% < y' < 0.7\%$ . For small  $\delta$  this implies  $y' \sim y_D \sim 1\%$  and  $x$  would be not much larger. However, for strong phase  $\delta \sim 20^\circ\text{--}50^\circ$ ,  $x_D$  could be several times larger than  $y_D \sim 1\%$ . With an active program at the B factories and CLEO-c, and the expectation that BESIII and LHCb would start running in 2008, it looks promising that  $x_D \sim 0.01$  to  $0.03$  can be discovered soon.

#### 4. DCPV in $B^+ \rightarrow J/\psi K^+$

One intriguing “prediction” we can make is  $\mathcal{A}_{J/\psi K^+} \neq 0$ .<sup>15</sup>

The  $B^+ \rightarrow J/\psi K^+$  is dominated by the color-suppressed  $b \rightarrow c\bar{c}s$  tree, while inclusion of the penguin in SM3 does not alter the weak phase, which is  $\simeq 0$ . But the full amplitude is likely carrying a strong phase  $\delta$ , since all color-suppressed modes observed




Fig. 3. (a)  $S_{J/\psi K^0}$  vs  $\phi_{sb}$ , (b)  $A_{J/\psi K^+}$  vs  $\delta$ , for  $m_{t'} = 300$  GeV and  $r_{sb} = 0.02, 0.025$  and  $0.03$ . Larger  $r_{sb}$  gives stronger variation.

so far seem enhanced, with effective underlying strong phase. Examples are  $B^0 \rightarrow D^0 \pi^0$ ,  $\pi^0 \pi^0$ . Although the strong phase in the latter is still not settled, the former has a strong phase  $\sim 30^\circ$  that is measured. The most relevant is  $B \rightarrow J/\psi K^*$ , where angular analysis gives strong phase difference between helicity amplitudes at order  $30^\circ$ .

The  $t'$  effect in the  $Z$  penguin brings the weak  $\phi_{sb}$  phase to  $P_{EW}$  amplitude. Unlike the above “hadronic” effects that enhance  $C$ , the virtual  $Z$  produces a small color-singlet  $c\bar{c}$  pair that exits without much interaction, thereby not accumulating much strong phase. While a little hand waving, we see that both weak and strong phases are present, the prerequisites for DCPV.

We plot  $S_{J/\psi K}$  vs  $\phi_{sb}$  in Fig. 3(a), for  $\delta = 0$ . Similar to  $\Delta S$ , which has  $S_{J/\psi K}$  as reference point,  $S_{J/\psi K}$  itself does dip downwards for  $\phi_{sb} \sim 65^\circ$ , reaching roughly 0.69. This does not change significantly when  $\delta$  remains small. In Fig. 3(b) we plot  $A_{J/\psi K^+}$  vs  $\delta$  for  $\phi_{sb} = 65^\circ$ . We find that<sup>15</sup>  $A_{J/\psi K^+}$  can reach above 1% for  $|\delta| \sim 30^\circ$ .

The experimental situation<sup>16</sup> is interesting. From  $A_{J/\psi K^+} \sim +0.03$  based on 89M  $B\bar{B}$ s, BaBar flipped sign by adding 35M, becoming  $-0.030 \pm 0.014 \pm 0.010$ , with larger

systematic error, and is now consistent with Belle value of  $-0.026 \pm 0.022 \pm 0.017$  based on 32M. The current world average is  $-0.024 \pm 0.014$ , based on 166M  $B\bar{B}$ s. But the world has now over 1000M  $B\bar{B}$ s and growing, thus, our 1% projection can be seriously probed. Note that the number could be higher,<sup>15</sup> e.g. in the less constrained  $Z'$  model. To realize a 1% measurement, it seems that one needs to work hard on systematic error. But this should be worthwhile if one wants to enter the “Super B factory” era, with 100 times more data, where any measurement of interest is likely to be systematics limited.

With luck, our prediction can be confirmed by 2008.

## Acknowledgments

I thank Makiko Nagashima and Andrea Soddu for collaboration.

## References

1. S. Giagu [CDF Collab.], hep-ex/0610044.
2. Plenary talk by M. Hazumi, this proceedings.
3. J. Erler, hep-ph/0604035.
4. A. Soddu in neutrino parallel session.
5. W.S. Hou, R.S. Willey and A. Soni, *Phys. Rev. Lett.* **58**, 1608 (1987).
6. A. Arhrib and W.S. Hou, *Eur. Phys. J.* **C27**, 555 (2003).
7. W.S. Hou, M. Nagashima and A. Soddu, hep-ph/0610385.
8. W.S. Hou, M. Nagashima and A. Soddu, *Phys. Rev. Lett.* **95**, 141601 (2005).
9. Plenary talk by R. Barlow, this proceedings.
10. W.S. Hou, M. Nagashima, G. Raz and A. Soddu, *JHEP* **0609**, 012 (2006).
11. R. Sinha, B. Misra and W.S. Hou, *Phys. Rev. Lett.* **97**, 131802 (2006).
12. W.S. Hou, H.n. Li, S. Mishima and M. Nagashima, hep-ph/0611107.
13. W.S. Hou, M. Nagashima and A. Soddu, *Phys. Rev.* **D72**, 115007 (2005).
14. Plenary talk by P. Pakhlov, this proceedings.
15. W.S. Hou, M. Nagashima and A. Soddu, hep-ph/0605080.
16. W.M. Yao *et al.* [Particle Data Group], *J. Phys.* **G33**, 1 (2006).